been identified to date (2). Apart from its substrate Rac and the regulators mentioned above, P-Rex1 has been shown to interact

STXB1 and DPYL2 in band 2, all migrating as expected for their MW, and STXB1 was also found in fraction 15, band 7. Other bands in fraction 3 were not analyzed, because careful evaluation of the surrounding lanes suggested that they likely contained proteins that bound nonspecifically to the beads. Norbin was recovered in fraction 4, band 3, and AP3B2 in fraction 11, band 5. Bands 4 and 6 contained only keratin and trypsin, so were not considered further.

Among the five proteins identified, we selected Norbin

P-Rex1 and Norbin interact directly and in cells

To confirm our initial identification of Norbin as a

vitro, ΔDEP*, and one mutant that couldnot,* ΔPH*. The EE-*ΔDEP *mutant coimmunoprecipitatedwith myc-Norbin as did full length EE-P-Rex1, whereas the EE-*ΔPH *mutant did not (Figure 4D). These results show that the PH domain of P-Rex1 is required for Norbin binding in vivo, suggesting that similar P-Rex1 domains confer the interaction with Norbin in vitro and in vivo.*

Norbin stimulates the Rac-GEF activity of P-Rex1 in vitro and in cells

To test whether the direct interaction between P-Rex1 and Norbin affects the Rac1-GEF activity of P-Rex1, purified GST-Norbin (as in Figures 3D and E, except not immobilized) was incorporated into a liposome-based in vitro Rac-GEF activity assay that measures the activation (GTPγS-loading) of Rac by P-Rex1, in the presence or absence of the P-Rex1 activators PIP$_3$ and Gβγ *(1,48). Here, we used Sf9-cell derived purified recombinant prenylated GDP-loaded EE-Rac1 and EE-*$G\beta_1\gamma_2$ *proteins, full-length EE-P-Rex1 (as in Figures 3D and E) and synthetic stearoyl-arachidonyl PIP*$_3$*. In the absence of EE-P-Rex1, 16% of Rac1 was active (GTPγS-loaded), and GST-Norbin had no effect on this basal activity, whereas EE-P-Rex1 increased Rac1 activity to 31%. The combined addition of EE-P-Rex1 and GST-Norbin at a molar ratio of 1:4 increased Rac1 activity to 44%, whereas GST again had no effect. These data show that GST-Norbin directly stimulates the basal Rac-GEF activity of P-Rex1. PIP*$_3$ and Gβγ proteins, also stimulated the *activity of P-Rex1, as expected, and GST-Norbin increased this further from 63 to 87% and from 71 to 91%, respectively, whereas GST again had no effect (Figure 5A). Therefore, Norbin directly stimulates the basal, PIP*$_3$- and Gβγ*-dependent Rac1-GEF activities of P-Rex1, modestly but significantly. The degree of P-Rex1 activation by Norbin was similar under all conditions, roughly additive to the effects of PIP*$_3$ and Gβγ*. This suggests that Norbin binding might increase the availability of P-Rex1 in the liposomes that contain* the prenylated Rac1 and Gβγ proteins and the PIP$_3$ *phospholipid, thereby bringing P-Rex1 into closer contact with its substrate and its other activators.*

To assess whether Norbin can also stimulate the Rac-GEF activity of P-Rex1 in vivo, we performed Pak-CRIB pull-down assays (49). HEK-293 cells overexpressing EE-P-Rex1 and/or myc-Norbin, or mock-transfected cells as controls, were serum-starved and stimulated with the GPCR ligand lysophosphatidic acid (LPA), which is known to stimulate GPCR-mediated P-Rex1 activation in HEK-293 cells (23), or they were mock-stimulated. Endogenous active (GTP-bound) Rac1 was isolated from total cell lysates using GST-Pak-CRIB immobilized on glutathione sepharose, and GTP-Rac1 and total Rac1 levels were analyzed by western blotting. In basal (serum-starved) cells, 0.40% of the total cellular Rac1 was active, and the expression of myc-Norbin significantly reduced this level of active Rac1 to 0.14%. As we aimed for a modest level of overexpression, EE-P-(L , a n d

In addition to stimulating the Rac-GEF activity of P-Rex1, Gβγ and PIP_3 also regulate the Rac-GEF by synergistically driving its translocation from the cytosol to the plasma membrane (26,27) To assess whether Norbin also affects the subcellular localization of P-Rex1, we carried out immunofluorescence microscopy in porcine aortic endothelial (PAE) cells. In addition, we assessed cell morphology. PAE cells have a kite-shaped morphology when basal, but adopt characteristic morphologies, namely lamellipodia formation, membrane ruffling and cell spreading, in response to P-Rex1-dependent activation of Rac1, through the induction of actomyosin cytoskeletal dynamics (1,50). PAE cell morphology thereby provides an indirect but sensitive readout of endogenous Rac1 activity that can be assessed on a cell-by-cell basis through imaging (2).

PAE cells expressing eGFP-P-Rex1 and/or myc-Norbin were serum-starved fixed, stained, and the subcellular localizations of eGFP-P-Rex1 and myc-Norbin as well as cell shape evaluated by widefield immunofluorescence imaging. The localization of eGFP-P-Rex1 appeared largely cytosolic when the Rac-GEF was expressed alone, although some cells showed a partial membrane localization of the GEF, with lamellipodia formation and membrane ruffling, as expected. myc-Norbin appeared also largely cytosolic when expressed alone, and these cells had a basal morphology. In contrast, coexpression of eGFP-P-Rex1 and myc-Norbin seemed to induce a translocation of both proteins to the plasma membrane, and stimulate lamellipodia, membrane ruffling and cell spreading (Figure 6A). To investigate if eGFP-P-Rex1 and myc-Norbin co-localize, we imaged these PAE cells using super-resolution structured illumination microscopy (SIM). This imaging confirmed that eGFP-P-Rex1 and myc-Norbin are largely localized in the cytoplasm, with eGFP-P-Rex1 showing a more homogenous distribution than myc-Norbin which was slightly more punctate. Some colocalzation of eGFP-P-Rex1 and myc-Norbin was seen in the cytoplasm. However, much more substantial colocalization of eGFP-P-Rex1 and myc-Norbin was observed in membrane ruffles at the cell edge (Figure 6B and Movie 1

2.3(i)-2 0.5c Rule9 (bi)6. std MP vie 1 << /MCID

cells that coexpressed eGFP-P-Rex1 and myc-Norbin,

membrane, respectively, we carried out subcellular fractionation of HEK293 cells expressing EE-P-Rex1 and/or myc-Norbin, using mock-transfected cells as a control. Total lysates of serum-starved cells were cleared of debris and nuclei by low-speed centrifugation, the post-nuclear supernatant fractionated further into membrane and cytosol fractions, and the amounts of P-Rex1 and Norbin protein in each fraction determined by western blotting with 6F12 P-Rex1 and Norbin C1 antibody (Figure 8A).

When EE-P-Rex1 was expressed alone, P-Rex1 protein was largely found in the cytosol fraction, as expected (1,26,27), and 11% of P-Rex1 from the post-nuclear supernatant were recovered in the membrane fraction. In contrast, coexpression with myc-Norbin decreased the amount of P-Rex1 protein in the cytosol fraction and increased its presence in the membrane fraction to 28%. (dh9 0 Tdr 0.01149 oTD [(m)1 0.8011 Tc 0.34599 Tw 82 cl Am) S m

phosphatidic acid might recruit Norbin to the plasma membrane upon cell stimulation. Furthermore, the N-terminus of Norbin can be palmitoylated, and this palmitoylation has been shown to recruit Norbin to early endosomes in hippocampal neurons (52). However, the plasma membrane localization of Norbin was not observed in that study, so palmitoylation is unlikely to be the mechanism of the plasma membrane recruitment we describe here. Finally, Norbin interacts directly and constitutively with many GPCRs (41,44,53), and also interacts in vivo with semaphorin 4C, a transmembrane protein that controls axon path finding and synapse structure (46). It is possible that P-Rex1 binding to Norbin increases the propensity of Norbin to bind to these plasma membrane localized proteins, and that P-Rex1 and Norbin are thus retained at the plasma membrane by multi-protein complex formation.

P7F1ix(v)1re Rac1

Norbin increased the Rac1-GEF activity of P-Rex1 directly in vitro, but its effect was minor compared to P-Rex1 activation by PIP_3 or Gβγ (1,18). In addition, the fold-stimulation of P-Rex1 activity by Norbin was similar, regardless of whether PIP_3 or Gβγ were present, which sugge

Norbin Stimulates the Catalytic Activity and Plasma Membrane Localization of the Guanine-Nucleotide Exchange Factor P-Rex1

Dingxin Pan, Mark A. Barber, Kirsti Hornigold, Martin J. Baker, Judit M. Toth, David Oxley and Heidi C. E. Welch

J. Biol. Chem. published online January 20, 2016

Access the most updated version of this article at doi: 10.1074/jbc.M115.686592

Alerts:
- When this article is cited
- When a correction for this article is posted

Click here to choose from all of JBC's e-mail alerts

Supplemental material:
http://www.jbc.org/content/suppl/2016/01/20/M115.686592.DC1.html

This article cites 0 references, 0 of which can be accessed free at
http://www.jbc.org/content/early/2016/01/20/jbc.M115.686592.full.html#ref-list-1

Downloaded from http://www.jbc.org/ at Ïã¸ÚÁù°ïÃÛÏ³ on January 25, 2016